

On the New Evaluation of an Old Integral

by William Walters and Michael Huber

ARL-TR-4689 December 2008

NOTICES

Disclaimers

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof.

Destroy this report when it is no longer needed. Do not return it to the originator.

Army Research Laboratory

Aberdeen Proving Ground, MD 21005-5069

ARL-TR-4689 December 2008

On the New Evaluation of an Old Integral

William Walters Weapons and Materials Research Directorate, ARL

> Michael Huber Muhlenberg College

Approved for public release; distribution is unlimited.

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)	2. REPORT TYPE	3. DATES COVERED (From - To)
December 2008	Final	July 2008–December 2008
4. TITLE AND SUBTITLE		5a. CONTRACT NUMBER
On the New Evaluation of an Old Integral		
		5b. GRANT NUMBER
		5c. PROGRAM ELEMENT NUMBER
6. AUTHOR(S)		5d. PROJECT NUMBER
William Walters and Michael Huber*		AH80
		5e. TASK NUMBER
		5f. WORK UNIT NUMBER
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory		8. PERFORMING ORGANIZATION REPORT NUMBER
ATTN: AMSRD-ARL-WM-7 Aberdeen Proving Ground, M	ГĊ	ARL-TR-4689
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)

12. DISTRIBUTION/AVAILABILITY STATEMENT

Approved for public release; distribution is unlimited.

13. SUPPLEMENTARY NOTES

*Muhlenberg College, Allentown, PA 18104

14. ABSTRACT

A simple technique is shown to evaluate $\int \frac{dx}{x^m (1+x^n)}$.

15. SUBJECT TERMS

calculus, integration, power series

16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON William Walters
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (Include area code)
UNCLASSIFIED	UNCLASSIFIED	UNCLASSIFIED	UL	12	410-278-6062

Standard Form 298 (Rev. 8/98)

Prescribed by ANSI Std. Z39.18

Contents

1.	Introduction	1
2.	Using Infinite Sums	1
3.	A Generalization	3
4.	Examples	3
Dis	stribution List	5

INTENTIONALLY LEFT BLANK.

1. Introduction

Consider the integral

$$\int \frac{dx}{x^m (1+x)}. (1)$$

In the *CRC Standard Mathematical Tables*, ¹ equation 1 can require repeated integral evaluations. Integral 87 shows

$$\int \frac{dx}{x^m (a+bx^n)^{p+1}} = \frac{1}{a} \int \frac{dx}{x^m (a+bx^n)^p} - \frac{b}{a} \int \frac{dx}{x^{m-n} (a+bx^n)^{p+1}}.$$
 (2)

Enter this integral into your favorite computer algebra system. Maple returns

$$\int \frac{dx}{x^m (1+x)} = \frac{x^{-m} (m-1)}{(1-m)m} + \frac{x^{-m} (m-1) \operatorname{LerchPhi} (-x, 1, -m)}{1-m},$$
(3)

which involves Lerch's Phi transcendent function, defined as

LerchPhi
$$(x, a, m) = \sum_{n=0}^{\infty} \frac{x^n}{(m+n)^a}$$
. (4)

Similarly, Mathematica gives

$$\int \frac{dx}{x^m (1+x)} = -\frac{x^{1-m} \text{ Hypergeometric } 2F1[1-m,1,2-m,x]}{-1+m} , \qquad (5)$$

which involves knowing the implementation of Hypergeometric functions. In this report, we seek to provide a simpler evaluation for integrals of this form (equation 1). We state up front that the exponent m need not be an integer.

2. Using Infinite Sums

The integrand of equation 1 can be rewritten as follows:

$$\frac{x^{-m}}{1+x}. (6)$$

By long division,

¹ Bever, W. H., Ed. *CRC Standard Mathematical Tables*. CRC Press, Inc.: West Palm Beach, FL, 1978.

$$\frac{x^{-m}}{1+x} = x^{-m} - x^{-m+1} + x^{-m+2} - x^{-m+3} + \cdots, \tag{7}$$

which is an infinite sum in the form

$$\sum_{i=0}^{\infty} (-1)^i x^{-m+i}.$$
(8)

Therefore, the integral becomes

$$\int \frac{dx}{x^{m} (1+x)} = \int \sum_{i=0}^{\infty} (-1)^{i} x^{-m+i} dx$$

$$= \sum_{i=0}^{\infty} \int (-1)^{i} x^{-m+i} dx$$

$$= \sum_{i=0}^{\infty} \frac{(-1)^{i} x^{-m+i+1}}{-m+i+1} + C, \tag{9}$$

for some constant C. Noting that

$$\frac{1}{1+x} = \sum_{i=0}^{\infty} (-1)^i x^i \tag{10}$$

converges for $x^2 < 1$ allows us to rewrite equation 9 as

$$\int \frac{dx}{x^m (1+x)} = \frac{1}{x^m} \sum_{i=0}^{\infty} \frac{(-1)^i x^{i+1}}{i-m+1} + C;$$
(11)

again, m need not be an integer. However, if m = 1, 2, 3, ..., then for $i = m-1, x^{-m+i} = x^{-1}$, so

$$\int \frac{dx}{x} = \ln x \tag{12}$$

for the i = m - 1 term, or

$$\int \frac{dx}{x^m (1+x)} = \frac{1}{x^m} \sum_{i=0}^{m-2} \frac{(-1)^i x^{i+1}}{i-m+1} + \ln x + \sum_{i=m}^{\infty} \frac{(-1)^i x^{i+1}}{i-m+1} + C.$$
 (13)

This allows us to sum past the singularity.

3. A Generalization

Generalizing, consider the following:

$$\int \frac{dx}{x^m (1+x^n)} = \int \frac{x^{-m} dx}{1+x^n}.$$
 (14)

From long division, the integrand can be rewritten as x^{-m} $(1 - x^n + x^{2n} - x^{3n} + ...)$,

or

$$\frac{x^{-m}}{1+x^n} = \sum_{i=0}^{\infty} (-1)^i x^{in-m}.$$
 (15)

Thus,

$$\int \frac{dx}{x^m \left(1 + x^n\right)} = \int \sum_{i=0}^{\infty} \left(-1\right)^i x^{in-m} dx = \sum_{i=0}^{\infty} \frac{\left(-1\right)^i x^{in-m+1}}{ni - m + 1} + C,$$
(16)

and we sum prior to integration past any singularity. As before, m and n need not be integers, and this technique converges for |x| < 1.

4. Examples

As a first example, we consider the case with m = 3, an integer. We start with equation 2 (shown previously), where m = 3, a = b = n = 1, and p = 0. Continuing, we find that

$$\int \frac{dx}{x^{3}(1+x)} = \int \frac{dx}{x^{3}} - \int \frac{dx}{x^{2}(1+x)}$$

$$= -\frac{x^{-2}}{2} - \left[\int \frac{dx}{x^{2}} - \int \frac{dx}{x(1+x)} \right]$$

$$= -\frac{x^{-2}}{2} + x^{-1} + \ln x - \ln (1+x) + C$$

$$= -\frac{1}{2x^{2}} + \frac{1}{x} + \ln x - \left[x - \frac{x^{2}}{2} + \frac{x^{3}}{3} - \frac{x^{4}}{4} + \cdots \right] + C$$

$$= -\frac{1}{2x^{2}} + \frac{1}{x} + \ln x - x + \frac{x^{2}}{2} - \frac{x^{3}}{3} + \frac{x^{4}}{4} - \cdots + C, \qquad (17)$$

which is valid for -1 < x < 1. Using our approach and equation 9,

$$\int \frac{dx}{x^3 (1+x)} = \int \sum_{i=0}^{\infty} (-1)^i x^{-3+i} dx$$

$$= \int \left[x^{-3} - x^{-2} + x^{-1} - x^0 + x^1 - \cdots \right] dx$$

$$= -\frac{1}{2x^2} + \frac{1}{x} + \ln x - x + \frac{x^2}{2} - \frac{x^3}{3} + \frac{x^4}{4} - \cdots + C, \tag{18}$$

which matches the CRC's solution. When m is a positive integer, equation 2 is adequate. However, as m gets large, the CRC's proposal will involve evaluating several integrals. Our technique is much simpler.

Now, suppose $m = \frac{1}{5}$, n = 3, and we keep a = b = 1 and p = 0. Equation 2 becomes

$$\int \frac{dx}{x^{\frac{1}{5}}(1+x^3)} = \int \frac{dx}{x^{\frac{1}{5}}} - \int \frac{dx}{x^{\frac{-14}{5}}(1+x^3)} \,. \tag{19}$$

While the first integral on the right-hand side is easy enough to evaluate, the second is not trivial, unless the process is repeated or the integrand is expanded in a power series. Turning to equation 15, we find

$$\int \frac{dx}{x^{\frac{1}{5}}(1+x^{3})} = \int \sum_{i=0}^{\infty} (-1)^{i} x^{3i-\frac{1}{5}} dx$$

$$= \int \left[x^{-\frac{1}{5}} - x^{\frac{14}{5}} + x^{\frac{29}{5}} - x^{\frac{44}{5}} + \cdots \right] dx$$

$$= \frac{5}{4} x^{\frac{4}{5}} - \frac{5}{19} x^{\frac{19}{5}} + \frac{5}{34} x^{\frac{34}{5}} - \frac{5}{49} x^{\frac{49}{5}} + \cdots + C. \tag{20}$$

We do not offer any practical applications for the integral. However, the proposed method converges for $x^2 < 1$ or when the independent variable is normalized, to guarantee that the normalized value of x is less than 1 when nondimensionalized; otherwise, the series diverges.

NO. OF

COPIES ORGANIZATION

1 DEFENSE TECHNICAL
(PDF INFORMATION CTR
only) DTIC OCA
8725 JOHN J KINGMAN RD
STE 0944
FORT BELVOIR VA 22060-6218

- 1 DIRECTOR
 US ARMY RESEARCH LAB
 IMNE ALC IMS
 2800 POWDER MILL RD
 ADELPHI MD 20783-1197
- 1 DIRECTOR
 US ARMY RESEARCH LAB
 AMSRD ARL CI OK TL
 2800 POWDER MILL RD
 ADELPHI MD 20783-1197
- DIRECTOR
 US ARMY RESEARCH LAB
 AMSRD ARL CI OK PE
 2800 POWDER MILL RD
 ADELPHI MD 20783-1197

ABERDEEN PROVING GROUND

1 DIR USARL AMSRD ARL CI OK TP (BLDG 4600)

NO. OF COPIES ORGANIZATION

- 1 DEPT OF AEROSPACE ENGR AND MECH UNIV OF ALABAMA S JONES PO BOX 870280 TUSCALOOSA AL 35487-0280
- 20 DEPT OF MATH
 AND CMPTR SCI
 MUHLENBERG COLLEGE
 M HUBER
 ALLENTOWN PA 18104
- 1 COMMANDER
 US ARMY RESEARCH OFFICE
 B LAMATTINA
 PO BOX 12211
 RTP NC 27709-2211
- 1 AIR FORCE ARMAMENT LAB AFATL DLJR D LAMBERT EGLIN AFB FL 32542-6810
- 2 DARPA W SNOWDEN S WAX 3701 N FAIRFAX DR ARLINGTON VA 22203-1714
- 1 LOS ALAMOS NATL LAB P HOWE MS P915 PO BOX 1663 LOS ALAMOS NM 87545
- 1 LOS ALAMOS NATL LAB L HULL MS A133 PO BOX 1663 LOS ALAMOS NM 87545
- 2 DIR LLNL D BAUM L099 M MURPHY PO BOX 808 MS L35 LIVERMORE CA 94550
- 2 SOUTHWEST RESEARCH INST C ANDERSON J WALKER PO DRAWER 28510 SAN ANTONIO TX 78228-0510

NO. OF COPIES ORGANIZATION

2 DETK R CICCARELLI W FLIS 3620 HORIZON DR KING OF PRUSSIA PA 19406

ABERDEEN PROVING GROUND

29 **DIR USARL** AMSRD ARL WM P PLOSTINS J SMITH AMSRD ARLWM BF D WEBB AMSRD ARL WM T P BAKER AMSRD ARL WM TC T FARRAND K KIMSEY L MAGNESS R MUDD D SCHEFFLER S SCHRAML S SEGLETES **B SORENSEN** R SUMMERS A TANK W WALTERS (10 CPS) **C WILLIAMS** AMSRD ARL WM TD T BJERKE M GREENFIELD E RAPACKI M RAFTENBERG